

# Challenge

# **EASY & EFFICIENT**

5 simple and fast experiments to measure the height of a building using a smartphone.



Discover The Smartphone Physics Challenge at [VULGARISATION.FR](http://VULGARISATION.FR)

«Physics Reimagined» team (Paris-Saclay University)



Precision: high



Difficulty: minimum

# Nº3. Free Fall Filmed


## Formula

$$H = \frac{1}{2} g t^2$$

## Material





1 ball



Sensor:  
**camera**

1 smartphone



Drop the ball from the top of the building. Film the fall and determine its duration.

$t$  = fall time of the ball,  
 $g = 9.8 \text{ ms}^{-2}$

*The formula does not consider air drag.*



Precision: high



Difficulty: minimum

# Nº21. Thales and the Shadows


## Formula

$$H = h \frac{l_2}{l_1}$$

## Material




1 tape measure



1 smartphone

Measure the shadow of a smartphone and the shadow of the building. Use Thales' method to determine the height of the building from the height of the smartphone.



$h$  = height of the smartphone  $l_2$  = shadow of the building,  $l_1$  = shadow of the smartphone



Precision: maximum



Difficulty: minimum

# Nº28. Picture with Scale

## Formula

$$H = \frac{d_2}{d_1} l$$

## Material





1 bar of known size



Sensor:  
camera

1 smartphone



Take a picture of the facade of the building, with the bar serving as a scale. Measure the sizes of the building and the bar on the picture.

$d_2$  = size of the building on the photo,  $d_1$  = size of the bar on the photo,  $l$  = actual size of the bar

*Minimize perspective distortion while taking the picture!*



Precision: maximum




Difficulty: minimum

# Nº35. Number of Steps

## Formula


$$H = Nh$$

## Material





Sensor:  
**accelerometer**

1 smartphone



Using the accelerometer, count the number of stair steps to the top of the building.

$N$  = number of steps,  
 $h$  = height of a step





Precision: intermediate



Difficulty: minimum

# Nº39. Acoustic Stopwatch


## Formula

$$H = v \frac{\delta t}{2}$$

## Material




2 balloons




Sensor:  
**microphone**

2 smartphones



Install an acoustic stopwatch application on both smartphones (Phyphox for example). Launch the application, a smartphone at the bottom of the building, one at the top. Trigger the timers by popping a balloon at the bottom, then stop the timers by popping a balloon at the top.



$v$  = speed of sound,  $\delta t$  = difference between the two chronometers

This project was imagined by Frédéric Bouquet (Paris-Saclay University) and Giovanni Organtini (Sapienza Università di Roma, Italy).

Physics: Frédéric Bouquet, Giovanni Organtini, Julien Bobroff

Videos, photos, gifs: Amel Kolli

Graphic design and illustrations:  
Anna Khazina

This project is a production of «Physics Reimagined» from Paris-Saclay University and CNRS. It benefited from the support of the IDEX Paris-Saclay and of the «Physique Autrement» Chair, held by the Paris-Sud Foundation and supported by the Air Liquide Group.