

C'EST

C'EST

C'EST

C'EST

QUOI

C'EST

QUOI

C'EST

QUOI

C'EST

QUOI

CETTE

C'EST

QUOI

CETTE

C'EST

QUOI

CETTE

C'EST

QUOI

CETTE

COURBE ?

C'EST

QUOI

CETTE

COURBE ?

C'EST

QUOI

CETTE

COURBE ?

**C'EST
QUOI
CETTE
COURBE ?**

C'EST

QUOI

CETTE

COURBE ?

**C'EST
QUOI
CETTE
COURBE ?**

**C'EST
QUOI
CETTE
COURBE ?**

Magnétorésistance

**C'EST
QUOI
CETTE
COURBE ?**

Magnéto-résistance VS

**C'EST
QUOI
CETTE
COURBE ?**

***Magnétorésistance VS
Champ***

**C'EST
QUOI
CETTE
COURBE ?**

***Magnétorésistance VS
Champ magnétique***

**C'EST
QUOI
CETTE
COURBE ?**

***Magnétorésistance VS
Champ magnétique***

**C'EST
QUOI
CETTE
COURBE ?**

***Magnétorésistance VS
Champ magnétique***

C'EST

QUOI

COURBÉ ?

CETTE

***Magnétorésistance VS
Champ magnétique***

C'EST

QUOI

**COUVRIR
CETTE**

***Magnétorésist
Champ magnét***

CE

QUOI

CETTE

COMI

Magnéto
Champ ma

3

qu

ETTE

88

Mag
Cham

TE

90

M
Ch





Albert



Albert Fert



Albert Fert Prix Nobel



**Albert Fert
Prix Nobel
2007**

Prix Nobel
2007
Albert Fert



Prix Nobel

Albert Fert

2007



x Nobel

Albert Fert

2007



Albert

2007

obel



AI

20




FIG. 2. Magnetoresistance of a $[(\text{Fe } 30 \text{ \AA})/(\text{Cr } 9 \text{ \AA})]_{40}$ superlattice of 4.2 K. The current is along $[110]$ and the field is in the layer plane along the current direction (curve *a*), in the layer plane perpendicular to the current (curve *b*), or perpendicular to the layer plane (curve *c*). The resistivity at zero field is $54 \mu\Omega \text{ cm}$. There is a small difference between the curves in increasing and decreasing field (hysteresis) that we have not represented in the figure. The superlattice is covered by a 100-\AA Ag protection layer. This means that the magnetoresistance of the superlattice alone should be slightly higher.

of Grünberg *et al.*⁴ and by the spin-polarized low-energy electron-diffraction experiments of Carbone and Alvarado.¹⁰ The AF coupling between the Fe layers has been ascribed to indirect exchange interactions through the Cr layers, but a theoretical model of these interactions is still lacking.^{4,9}

The magnetoresistance of the Fe/Cr superlattices has been studied by a classical ac technique on small rectangular samples. Examples of magnetoresistance curves at 4.2 K are shown in Figs. 2 and 3. The resistance decreases during the magnetization process and becomes practically constant when the magnetization is saturated. The curves *a* and *b* in Fig. 2 are obtained for applied fields in the plane of layers in the longitudinal and transverse directions, respectively. The field H_S is the field needed to overcome the AF couplings and to saturate the magnetization (compare with Fig. 1). In contrast, fields applied perpendicularly to the layers (curve *c*) have to overcome not only the AF coupling but also the magnetic anisotropy, so that the magnetoresistance is saturated at a field higher than H_S .

The most remarkable result exhibited in Figs. 2 and 3 is the huge value of the magnetoresistance. For $t_{\text{Cr}}=9 \text{ \AA}$ and $T=4.2 \text{ K}$, see Fig. 2, there is almost a factor of 2 between the resistivities at zero field and in the saturated state, respectively (in absolute value, the resistivity change is about $23 \mu\Omega \text{ cm}$). By comparison of the results for three different samples in Fig. 3, it can be seen




FIG. 3. Magnetoresistance of three Fe/Cr superlattices at 4.2 K. The current and the applied field are along the same $[110]$ axis in the plane of the layers.




FIG. 2. Magnetoresistance of a $[(\text{Fe } 30 \text{ \AA})/(\text{Cr } 9 \text{ \AA})]_{40}$ superlattice of 4.2 K. The current is along $[110]$ and the field is in the layer plane along the current direction (curve *a*), in the layer plane perpendicular to the current (curve *b*), or perpendicular to the layer plane (curve *c*). The resistivity at zero field is $54 \mu\Omega \text{ cm}$. There is a small difference between the curves in increasing and decreasing field (hysteresis) that we have not represented in the figure. The superlattice is covered by a 100-\AA Ag protection layer. This means that the magnetoresistance of the superlattice alone should be slightly higher.

of Grünberg *et al.*⁴ and by the spin-polarized low-energy electron-diffraction experiments of Carbone and Alvarado.¹⁰ The AF coupling between the Fe layers has been ascribed to indirect exchange interactions through the Cr layers, but a theoretical model of these interactions is still lacking.^{4,9}

The magnetoresistance of the Fe/Cr superlattices has been studied by a classical ac technique on small rectangular samples. Examples of magnetoresistance curves at 4.2 K are shown in Figs. 2 and 3. The resistance decreases during the magnetization process and becomes practically constant when the magnetization is saturated. The curves *a* and *b* in Fig. 2 are obtained for applied fields in the plane of layers in the longitudinal and transverse directions, respectively. The field H_S is the field needed to overcome the AF couplings and to saturate the magnetization (compare with Fig. 1). In contrast, fields applied perpendicularly to the layers (curve *c*) have to overcome not only the AF coupling but also the magnetic anisotropy, so that the magnetoresistance is saturated at a field higher than H_S .

The most remarkable result exhibited in Figs. 2 and 3 is the huge value of the magnetoresistance. For $t_{\text{Cr}}=9 \text{ \AA}$ and $T=4.2 \text{ K}$, see Fig. 2, there is almost a factor of 2 between the resistivities at zero field and in the saturated state, respectively (in absolute value, the resistivity change is about $23 \mu\Omega \text{ cm}$). By comparison of the results for three different samples in Fig. 3, it can be seen




FIG. 3. Magnetoresistance of three Fe/Cr superlattices at 4.2 K. The current and the applied field are along the same $[110]$ axis in the plane of the layers.




FIG. 2. Magnetoresistance of a $[(\text{Fe } 30 \text{ \AA})/(\text{Cr } 9 \text{ \AA})]_{40}$ superlattice of 4.2 K. The current is along $[110]$ and the field is in the layer plane along the current direction (curve *a*), in the layer plane perpendicular to the current (curve *b*), or perpendicular to the layer plane (curve *c*). The resistivity at zero field is $54 \mu\Omega \text{ cm}$. There is a small difference between the curves in increasing and decreasing field (hysteresis) that we have not represented in the figure. The superlattice is covered by a 100-\AA Ag protection layer. This means that the magnetoresistance of the superlattice alone should be slightly higher.

of Grünberg *et al.*⁴ and by the spin-polarized low-energy electron-diffraction experiments of Carbone and Alvarado.¹⁰ The AF coupling between the Fe layers has been ascribed to indirect exchange interactions through the Cr layers, but a theoretical model of these interactions is still lacking.^{4,9}

The magnetoresistance of the Fe/Cr superlattices has been studied by a classical ac technique on small rectangular samples. Examples of magnetoresistance curves at 4.2 K are shown in Figs. 2 and 3. The resistance decreases during the magnetization process and becomes practically constant when the magnetization is saturated. The curves *a* and *b* in Fig. 2 are obtained for applied fields in the plane of layers in the longitudinal and transverse directions, respectively. The field H_S is the field needed to overcome the AF couplings and to saturate the magnetization (compare with Fig. 1). In contrast, fields applied perpendicularly to the layers (curve *c*) have to overcome not only the AF coupling but also the magnetic anisotropy, so that the magnetoresistance is saturated at a field higher than H_S .

The most remarkable result exhibited in Figs. 2 and 3 is the huge value of the magnetoresistance. For $t_{\text{Cr}}=9 \text{ \AA}$ and $T=4.2 \text{ K}$, see Fig. 2, there is almost a factor of 2 between the resistivities at zero field and in the saturated state, respectively (in absolute value, the resistivity change is about $23 \mu\Omega \text{ cm}$). By comparison of the results for three different samples in Fig. 3, it can be seen









FIG. 3. Magnetoresistance of three Fe/Cr superlattices at 4.2 K. The current and the applied field are along the same $[110]$ axis in the plane of the layers.






















R




H

R




H

R




H

R




H

R



H



R



H

R



H



R



H



R



H



R



H



R




H



R




H







R






H








































































































































































































































































- ↑
- ↑
- ↑
- ↑
- ↓
- ↓
- ↓
- ↓


























R


















R






H








H



H







R







H





































































































































R









H




























































R





H










R



H





R




H



R




H







R




H






R




H










R








H












































R




H












R






H

































































FIG. 2. Magnetoconductance of a $[(\text{Fe } 30 \text{ \AA})/(\text{Cr } 9 \text{ \AA})]_{40}$ superlattice of 4.2 K. The current is along $[110]$ and the field is in the layer plane along the current direction (curve *a*), in the layer plane perpendicular to the current (curve *b*), or perpendicular to the layer plane (curve *c*). The resistivity at zero field is $54 \mu\Omega \text{ cm}$. There is a small difference between the curves in increasing and decreasing field (hysteresis) that we have not represented in the figure. The superlattice is covered by a 100-\AA Ag protection layer. This means that the magnetoconductance of the superlattice alone should be slightly higher.

of Grünberg *et al.*⁴ and by the spin-polarized low-energy electron-diffraction experiments of Carbone and Alvarado.¹⁰ The AF coupling between the Fe layers has been ascribed to indirect exchange interactions through the Cr layers, but a theoretical model of these interactions is still lacking.^{4,9}

The magnetoconductance of the Fe/Cr superlattices has been studied by a classical ac technique on small rectangular samples. Examples of magnetoconductance curves at 4.2 K are shown in Figs. 2 and 3. The resistance decreases during the magnetization process and becomes practically constant when the magnetization is saturated. The curves *a* and *b* in Fig. 2 are obtained for applied fields in the plane of layers in the longitudinal and transverse directions, respectively. The field H_S is the field needed to overcome the AF couplings and to saturate the magnetization (compare with Fig. 1). In contrast, fields applied perpendicularly to the layers (curve *c*) have to overcome not only the AF coupling but also the magnetic anisotropy, so that the magnetoconductance is saturated at a field higher than H_S .

The most remarkable result exhibited in Figs. 2 and 3 is the huge value of the magnetoconductance. For $t_{\text{Cr}}=9 \text{ \AA}$ and $T=4.2 \text{ K}$, see Fig. 2, there is almost a factor of 2 between the resistivities at zero field and in the saturated state, respectively (in absolute value, the resistivity change is about $23 \mu\Omega \text{ cm}$). By comparison of the results for three different samples in Fig. 3, it can be seen





FIG. 3. Magnetoconductance of three Fe/Cr superlattices at 4.2 K. The current and the applied field are along the same $[110]$ axis in the plane of the layers.


LE

LE GRAPHISME

LE GRAPHISME




LE GRAPHISME




Chloé

LE GRAPHISME



**Chloé
PASSAVANT**


LE GRAPHISME



**Chloé
PASSAVANT**

DSAA


LE GRAPHISME



**Chloé
PASSAVANT**

DSAA D


LE GRAPHISME



**Chloé
PASSAVANT**

DSAA DI


LE GRAPHISME



**Chloé
PASSAVANT**

DSAA DIS


LE GRAPHISME



**Chloé
PASSAVANT**

**DSAA DIS
Ecole**


LE GRAPHISME



**Chloé
PASSAVANT**

**DSAA DIS
Ecole Estienne**


LE GRAPHISME



**Chloé
PASSAVANT**

**DSAA DIS
Ecole Estienne**

LE GRAPHISME




**Chloé
PASSAVANT**

**DSAA DIS
Ecole Estienne**

LE GRAPHISME


**Chloé
PASSAVANT**



**DSAA DIS
Ecole Estienne**

LE GRAPHISME

**Chloé
PASSAVANT**




**DSAA DIS
Ecole Estienne**

GRAPHISME

**Chloé
PASSAVAN**


**DSAA DIS
Ecole Estienne**



RAPHISME

**Chloé
PASSAV**


**SAA DIS
cole Estienne**



PHISME

**Chloé
PASSI**


**DIS
Estienne**



ISME

**Chl
PAS**

ienne



ME

C
E


e




LA

LA PHYSIQUE

LA PHYSIQUE




LA PHYSIQUE




Julien

LA PHYSIQUE




Julien BOBROFF

LA PHYSIQUE




Julien BOBROFF

LA PHYSIQUE




LA PHYSIQUE

Frederic




LA PHYSIQUE

Frederic BOUQUET




LA PHYSIQUE

Frederic BOUQUET




LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET

La




LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET

La Physique




LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET


La Physique Autrement



LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET




La Physique Autrement
Laboratoire

LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET




La Physique Autrement
Laboratoire de

LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET




La Physique Autrement
Laboratoire de Physique

LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET




La Physique Autrement
Laboratoire de Physique des

LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET




La Physique Autrement
Laboratoire de Physique des Solides

LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET




La Physique Autrement
Laboratoire de Physique des Solides
Université

LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET




La Physique Autrement
Laboratoire de Physique des Solides
Université Paris

LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET




La Physique Autrement
Laboratoire de Physique des Solides
Université Paris-

LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET




La Physique Autrement
Laboratoire de Physique des Solides
Université Paris-Sud,

LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET




La Physique Autrement
Laboratoire de Physique des Solides
Université Paris-Sud, CNRS

LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET




La Physique Autrement
Laboratoire de Physique des Solides
Université Paris-Sud, CNRS

LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUET

La Physique Autrement
Laboratoire de Physique des Solides
Université Paris-Sud, CNRS




LA PHYSIQUE




Julien BOBROFF
Frederic BOUQUIN

Physique Autrement
Institut de Physique des Solides
Université Paris-Sud, CNRS




LA PH




Julien BOBR
Frederic BOU

ique Autrement
e de Physique des Solides
Paris-Sud, CNRS



LA PH




Julien BO
Frederic P

Autrement

Physique des Solides
UMR 5086, CNRS



LA




**Julien
Freder**

ement

les Solides

S

7



Julia
Frederick

t
les



W

WW

WWW

WWW.

WWW . VULGARISATION

WWW . VULGARISATION .

WWW . VULGARISATION . FR

WWW . VULGARISATION . FR



WWW . VULGARISATION . FR



WWW . VULGARISATION . FR

